## SOLUTION TO PROBLEM 12229

## TOMMASO MANNELLI MAZZOLI

Proposed by M. Omarjee (France).

Let  $f: [0,1] \to \mathbb{R}$  be a function that has continuous derivatives and that satisfies f(0) = f(1) and  $\int_0^1 f(x) dx = 0$ . Prove

$$30240 \left( \int_0^1 x f(x) \, \mathrm{d}x \right)^2 \le \int_0^1 \left( f''(x) \right)^2 \, \mathrm{d}x.$$

Solution. By the Cauchy-Schwarz inequality

$$\left(\int_0^1 f''(x)g(x) \, \mathrm{d}x\right)^2 \le \int_0^1 (f''(x))^2 \, \mathrm{d}x \cdot \int_0^1 g^2(x) \, \mathrm{d}x \quad \text{for every } g \in \mathrm{L}^2(0,1).$$

After applying integration by parts twice, we find

$$\int_0^1 f''(x)g(x) \, \mathrm{d}x = g(x)f'(x)\Big|_0^1 - \int_0^1 g'(x)f'(x) \, \mathrm{d}x$$

$$= g(x)f'(x)\Big|_0^1 - g'(x)f(x)\Big|_0^1 + \int_0^1 g''(x)f(x) \, \mathrm{d}x$$

$$= g(1)f'(1) - g(0)f'(0) - f(0)[g'(1) - g'(0)] + \int_0^1 g''(x)f(x) \, \mathrm{d}x.$$

Let

$$g(x) = \frac{1}{6}x(x-1)\left(x-\frac{1}{2}\right) = \frac{1}{12}\left(2x^3 - 3x^2 + x\right).$$

We have that:

- g(0) = g(1) = 0;
- g'(0) = g'(1);
- g''(x) = x for all  $x \in [0, 1]$ .

Thus,

$$\int_0^1 f''(x)g(x) \, \mathrm{d}x = \int_0^1 x f(x) \, \mathrm{d}x.$$

Then

$$\int_0^1 g^2(x) \, dx = \frac{1}{36} \int_0^1 x^2 (x - 1)^2 \left( x - \frac{1}{2} \right)^2 dx =$$

$$= \frac{1}{36} \int_0^1 \left( x^6 - 3x^5 + \frac{13x^4}{4} - \frac{3x^3}{2} + \frac{x^2}{4} \right) dx = \frac{1}{36} \cdot \frac{1}{840} = \frac{1}{30240}.$$

Therefore,

$$30240 \left( \int_0^1 x f(x) \, \mathrm{d}x \right)^2 \le \int_0^1 (f''(x))^2 \, \mathrm{d}x.$$